

CHARACTER FORMULAS ON COHOMOLOGY OF DEFORMATIONS OF HILBERT SCHEMES OF $K3$ SURFACES

LETAO ZHANG

ABSTRACT. Let X be a Kähler deformation of the Hilbert scheme of n points on a $K3$ surface. We compute the graded character formula of the generic Mumford-Tate group representation on the cohomology ring of X . Also, we derive a generating series for deducing the number of canonical Hodge classes of degree $2n$.

CONTENTS

1. Introduction	1
2. Cohomology of Hilbert Schemes of Points on $K3$ surfaces	2
3. Decomposition of Cohomology Ring	4
4. Generating Series of the Character of the Middle Cohomology $H^{2n}(X, \mathbb{Q})$	12
5. Appendix: Table of Nonstable G_X Representations	15
References	16

1. INTRODUCTION

Let S be a $K3$ surface, $S^{[n]}$ be the Hilbert scheme of n points on S , and X be a Kähler deformation of $S^{[n]}$; each point in $S^{[n]}$ corresponds to a subscheme of S whose Hilbert polynomial is the constant n .

Denote by G_X the Mumford-Tate group of X ; the invariants of the G_X action on $H^*(X, \mathbb{Q})$ correspond to the *canonical Hodge classes*, which are Hodge classes that remain Hodge under deformations. Chern classes of the tangent bundle are examples of such canonical Hodge classes, but generally additional Hodge classes will exist, since the dimension of the G_X invariants on $H^{2n}(X, \mathbb{Q})$ is greater than the number of monomials in the Chern classes. As for the action of G_X , we can consider the character of the representation on the middle cohomology for each n .

Theorem 1. *Let $M(q) := \sum_{n=0}^{\infty} \text{Char}(H^{2n}(X, \mathbb{Q})) \cdot q^n$ be the generating series of the character of the representation on the middle cohomology of X . Then*

$$M(q) = \left(1 + \sum_{k=1}^{\infty} 2(-1)^k q^{\frac{k(k+1)}{2}} \right) \left(\prod_{m=1}^{\infty} \det(I_{24} - gq^m) \right)^{-1},$$

where $g \in G_X$ is diagonalizable, I_N is a $N \times N$ identity matrix, and $\det(I_{24} - gt^m) = (1 - z^{-2}t^m) \cdot (1 - z^2t^m) \det(I_{22} - g|_{H^2(S, \mathbb{Q})}t^m)$.

Let $l \in H_2(X, \mathbb{Z})$ be a line class in $\mathbb{P}^n \subset X$. Hassett and Tschinkel conjecture in [9] that $(l, l) = -\frac{n+3}{2}$ and show that $(l, l) = -\frac{5}{2}$ for the case where $n = 2$ in [8]. For $n = 3$, Harvey, Hassett and Tschinkel [7] show that $(l, l) = -3$ and give a concrete expression for the Lagrangian hyperplane class. For the case of $n = 4$, Bakker and Jorza [1] show that $(l, l) = -\frac{7}{2}$, and also give an expression $[\mathbb{P}^4]$. However, it is more difficult to compute the class $[\mathbb{P}^n]$ for larger n . One possible approach to proving the conjecture is to find all the canonical Hodge classes in the middle cohomology of X for each n ; future work could provide possible candidates for the class of $[\mathbb{P}^n]$ in terms of the line class. As for the ring structure, Verbitsky [15] shows that there is an embedding $\text{Sym}^n H^2(S, \mathbb{Q}) \hookrightarrow H^{2n}(S^{[n]}, \mathbb{Q})$, but much about the ring structure of $H^*(X, \mathbb{Q})^{G_X}$ is still unknown, e.g. relations in the subalgebra generated by $H^*(S, \mathbb{Q})$ for each $H^*(X, \mathbb{Q})^{G_X}$.

Acknowledgements. I am very grateful to my advisor Brendan Hassett for introducing me to this problem, and for his warm support and encouragement. Thanks are also due to Lothar Göttsche, Radu Laza, and Anthony Várilly-Alvarado for interesting discussions and insightful remarks. I also appreciate Eyal Markman's illuminating questions which may lead to future research topics.

2. COHOMOLOGY OF HILBERT SCHEMES OF POINTS ON $K3$ SURFACES

In this section, we review some classical results about $S^{[n]}$. Let $\langle \cdot, \cdot \rangle$ denote the Beauville-Bogomolov form on the cohomology group $H^2(S^{[n]}, \mathbb{Z})$. Since S is a $K3$ surface, $H^2(S^{[n]}, \mathbb{Z})_{\langle \cdot, \cdot \rangle}$ ($n \geq 2$) is the direct sum given by Beauville in [2] by

$$(2.1) \quad H^2(S^{[n]}, \mathbb{Z})_{\langle \cdot, \cdot \rangle} = H^2(S, \mathbb{Z})_{\langle \cdot, \cdot \rangle} \oplus_{\perp} \mathbb{Z}\delta, \quad (\delta, \delta) = -2(n-1),$$

where $(\cdot, \cdot)$ is the intersection form on $H^2(S, \mathbb{Z})$, and 2δ is the class of the corresponding big diagonal divisor $\Delta^{[n]} \subset S^{[n]}$ parameterizing nonreduced subschemes.

In [14], Nakajima constructs generators for the cohomology ring of Hilbert schemes of points of any projective surface. Lehn and Sorger [12] then show how $H^*(S, \mathbb{Q})$ generates $H^*(S^{[n]}, \mathbb{Q})$ as a graded ring.

Let $A = H^*(S, \mathbb{Q})[2]$ denote the shifted cohomology ring weighted by $-2, 0, 2$. Correspondingly, let $\mathbb{H}_n = H^*(S^{[n]}, \mathbb{Q})[2n]$ denote the shifted cohomology ring weighted by $-2n, \dots, 2n$. Note that the weight shifting here is not the Tate twist notation for Hodge classes. Define a linear form T on A by $T(a) := -\int_{[S]} a$, and on $\mathbb{H}_n$ by $T(a) := (-1)^n \int_{S^{[n]}} a$. Now consider $A^{\otimes n}$, with the usual product given by

$$(a_1 \otimes \cdots \otimes a_n) \cdot (b_1 \otimes \cdots \otimes b_n) = (a_1 b_1) \otimes \cdots \otimes (a_n b_n) ,$$

T extends to $A^{\otimes n}$ via

$$T(a_1 \otimes \cdots \otimes a_n) = T(a_1) \cdots T(a_n)$$

We also have the symmetric group $\mathfrak{S}_n$ action on the n -fold tensor given by

$$\pi(a_1 \otimes \cdots \otimes a_n) = a_{\pi^{-1}(1)} \otimes \cdots \otimes a_{\pi^{-1}(n)} .$$

For any partition $n = n_1 + \cdots + n_k$, we have a homomorphism

$$A^{\otimes n} \rightarrow A^{\otimes k}$$

$$a_1 \otimes \cdots \otimes a_n \mapsto (a_1 \cdots a_{n_1}) \otimes \cdots \otimes (a_{n_1+\cdots+n_{k-1}+1} \cdots a_n)$$

Given a finite set I with n elements, let $\{A_i\}_{i \in I}$ be a family of copies of A indexed by I . Let $[n]$ denote $\{1, \dots, n\}$; we define

$$A^{\otimes I} := \left(\bigoplus_{f: [n] \xrightarrow{\cong} I} A_{f(1) \otimes \cdots \otimes f(n)} \right) / \mathfrak{S}_n$$

Finally, given a surjection $\phi : I \rightarrow J$ between two index sets, there is an induced multiplication

$$\phi^* : A^{\otimes I} \rightarrow A^{\otimes J} ,$$

and let

$$\phi_* : A^{\otimes J} \rightarrow A^{\otimes I}$$

be the adjoint of ϕ^* , satisfying

$$\phi_*(a \cdot \phi^*(b)) = \phi_*(a) \cdot b ,$$

where $a \in A^{\otimes I}$ and $b \in A^{\otimes J}$.

Denote by $\langle \pi \rangle \backslash [n]$ the set of orbits of $[n]$ under the action of π . Define

$$A\{\mathfrak{S}_n\} := \bigoplus_{\pi \in \mathfrak{S}_n} A^{\otimes \langle \pi \rangle \backslash [n]} \cdot \pi$$

$A\{\mathfrak{S}_n\}$ admits an action of $\sigma \in \mathfrak{S}_n$, induced by the bijection

$$\sigma : \langle \pi \rangle \setminus [n] \rightarrow \langle \sigma \pi \sigma^{-1} \rangle \setminus [n], \quad x \mapsto \sigma x.$$

This gives an automorphism of $A\{\mathfrak{S}_n\}$ given by

$$\tilde{\sigma} : a \cdot \pi \mapsto \sigma^*(\sigma \pi \sigma^{-1}).$$

Denote by $A^{[n]}$ the invariants under this action, then we have the graded isomorphism between the vector spaces [12]

$$A^{[n]} = \sum_{\|\alpha\|=n} \bigotimes_i \text{Sym}^{\alpha_i} A$$

where $\alpha = (1^{\alpha_1}, 2^{\alpha_2}, \dots, n^{\alpha_n})$ runs all partitions of n . For the case of $K3$ surfaces, Lehn and Sorger prove

Theorem 2. [12] *There is a canonical isomorphism of graded rings*

$$(H^*(S, \mathbb{Q})[2])^{[n]} \xrightarrow{\cong} H^*(S^{[n]}, \mathbb{Q})[2n].$$

3. DECOMPOSITION OF COHOMOLOGY RING

In this section, our goal is to decompose $H^*(X, \mathbb{Q})$ into irreducibles using representation theory.

3.1. Characters of Representations. We summarize general results on representations of complex (or split) orthogonal groups from [4].

Let $\mathfrak{g}$ be a semisimple Lie algebra, Λ be its weight lattice, and $\mathbb{Z}[\Lambda]$ be the integral group ring of the abelian group Λ . For each weight $\lambda \in \Lambda$, let $e(\lambda)$ denote the basis element in $\mathbb{Z}[\Lambda]$, so that each element in $\mathbb{Z}[\Lambda]$ can be written as the finite sum $\sum_{\lambda} n_{\lambda} \cdot e(\lambda)$. Denote by $R(\mathfrak{g})$ the ring of isomorphism classes of finite-dimensional representations associated to $\mathfrak{g}$. For each class $[V]$, $[V] = [V'] + [V'']$ whenever $V = V' \oplus V''$, and the product of two classes is defined as $[V] \cdot [W] = [V \otimes W]$. Define the character homomorphism

$$\text{Char} : R(\mathfrak{g}) \rightarrow \mathbb{Z}[\Lambda]$$

by $\text{Char}[V] = \sum \#(V_{\lambda}) \cdot e(\lambda)$, where V_{λ} is the weight space of V for the weight λ and $\#(V_{\lambda})$ is the multiplicity of V_{λ} in V . The Weyl group $\mathfrak{W}$ acts on $\mathbb{Z}[\Lambda]$ and the image of Char is contained in the ring of invariants $\mathbb{Z}[\Lambda]^{\mathfrak{W}}$.

Let $\omega_1, \dots, \omega_n$ be fundamental weights of $\mathfrak{g}$. Recall that fundamental weights have the property that any highest weight may be expressed uniquely as a nonnegative integral linear combination of them; they are free generators for the lattice Λ . Let Γ_i ($i = 1, \dots, n$) be the classes in $R(\mathfrak{g})$ of the irreducible representations of highest weight ω_i ($i = 1, \dots, n$). We have the following theorem.

Theorem 3. [4] *The representation ring $R(\mathfrak{g})$ is a polynomial ring on the variables $\Gamma_1, \dots, \Gamma_n$, and the homomorphism $\text{Char} : R(\mathfrak{g}) \rightarrow \mathbb{Z}[\Lambda]^{\mathfrak{W}}$ is an isomorphism.*

Thus decomposing V into irreducible $\mathfrak{g}$ representations is equivalent to finding its character polynomial.

Example 1. [4] Let $\mathfrak{g} = \mathfrak{so}_{2n}\mathbb{C}$ and $V \cong \mathbb{C}^{2n}$ be its standard representation. Its weight lattice Λ is $\text{span}\{L_1, \dots, L_n, (\sum L_i)/2\}$ (see Lecture 19 in [4] for detailed explanation). For $\mathfrak{so}_{2n}\mathbb{C}$, fundamental weights are $L_1, L_1 + L_2, \dots, L_1 + \dots + L_{n-1}, (L_1 + \dots + L_n)/2, (L_1 + \dots + L_{n-1} - L_n)/2$

corresponding to irreducible representations $V, \bigwedge^2 V, \dots, \bigwedge^{n-1} V$ and the half-spin representations S^+ and S^- . Set $t_i = e(L_i)$, $t_i^{-1} = e(-L_i)$, $t_i^{+1/2} = e(L_i/2)$, $t_i^{-1/2} = e(-L_i/2)$, $\text{Char}(\bigwedge^k V)$ is the k th elementary symmetric polynomial, denoted by D_k , of the $2n$ elements $t_1, t_1^{-1}, \dots, t_n, t_n^{-1}$. The character D^+ (resp. D^-) of S^+ (resp. S^-) is the sum $\sum t_1^{\pm 1/2} \dots t_n^{\pm 1/2}$, where the number of plus signs is even (resp. odd). Thus,

$$R(\mathfrak{so}_{2n}\mathbb{C}) = \mathbb{Z}[\Lambda]^{\mathfrak{W}} = \mathbb{Z}[D_1, \dots, D_{n-2}, D^+, D^-]$$

Example 2. [4] In the case of $\mathfrak{g} = \mathfrak{so}_{2n+1}\mathbb{C}$, its standard representation is $V \cong \mathbb{C}^{2n+1}$ and its weight lattice is the same as $\mathfrak{so}_{2n}\mathbb{C}$. But the fundamental weights are

$$L_1, L_1 + L_2, \dots, L_1 + L_2 + \dots + L_{n-1}, (L_1 + \dots + L_n)/2$$

corresponding to irreducible representations $V, \bigwedge^2 V, \dots, \bigwedge^{n-1} V$ and the spin representation S . $\text{Char}(\bigwedge^k V)$ here is the k th elementary symmetric polynomial, denoted by B_k , of $2n+1$ elements $t_1, t_1^{-1}, \dots, t_n, t_n^{-1}$ and 1. Denote by B the character of S , which is the n th symmetric polynomial in variables $t_i^{\frac{1}{2}} + t_i^{-\frac{1}{2}}$. By applying Theorem 3 we obtain

$$R(\mathfrak{so}_{2n+1}\mathbb{C}) = \mathbb{Z}[\Lambda]^{\mathfrak{W}} = \mathbb{Z}[B_1, \dots, B_{n-1}, B]$$

In general, we have $\mathfrak{so}_{2n}\mathbb{C} \subset \mathfrak{so}_{2n+1}\mathbb{C}$. If Γ_λ is the irreducible $\mathfrak{so}_{2n+1}\mathbb{C}$ representation of highest weight $\lambda = (\lambda_1 \geq \dots \geq \lambda_n \geq 0)$, the restriction representation is

$$\text{Res}_{\mathfrak{so}_{2n}\mathbb{C}}^{\mathfrak{so}_{2n+1}\mathbb{C}} \Gamma_\lambda = \oplus_{\bar{\lambda}} \Gamma_{\bar{\lambda}}$$

where $\bar{\lambda} = (\bar{\lambda}_1, \dots, \bar{\lambda}_n)$ satisfies

$$\lambda_1 \geq \bar{\lambda}_1 \geq \lambda_2 \geq \bar{\lambda}_2 \geq \dots \geq \bar{\lambda}_{n-1} \geq \lambda_n \geq |\bar{\lambda}_n|,$$

and the $\bar{\lambda}_i$ and λ_i are either all integers or all half integers.

Given a finite dimensional $\mathfrak{so}_{2n+1}\mathbb{C}$ representation W , if it is induced by the inclusion $\mathfrak{so}_{2n}\mathbb{C} \subset \mathfrak{so}_{2n+1}\mathbb{C}$ and all the weights of $\mathfrak{so}_{2n}\mathbb{C}$ representation are integer-valued, then so are the weights of the $\mathfrak{so}_{2n+1}\mathbb{C}$ representation. This implies that the character of the $\mathfrak{so}_{2n+1}\mathbb{C}$ representation will be in $\mathbb{Z}[B_1, \dots, B_{n-1}]$.

3.2. Group actions on cohomologies. Let V be a $\mathbb{Q}$ vector space, and let $\mathbb{S}(\mathbb{R})$ denote $\mathbb{C}^*$, regarded as a Lie group.

$$S^1 = \{z \in \mathbb{C}^* : z\bar{z} = 1\}$$

is a maximal compact subgroup of $\mathbb{S}(\mathbb{R})$.

Definition 3.3. ([6], I.A) A *Hodge structure* of weight n is given by a representation on $V_{\mathbb{R}} := V \otimes_{\mathbb{Q}} \mathbb{R}$

$$\tilde{\varphi} : \mathbb{S}(\mathbb{R}) \rightarrow \mathrm{GL}(V_{\mathbb{R}})$$

such that for $r \in \mathbb{R}^* \subset \mathbb{S}(\mathbb{R})$, $\tilde{\varphi}(r) = r^n \mathrm{id}_V$.

We can see that $\varphi := \tilde{\varphi}|_{S^1}$ gives a representation

$$\varphi : S^1 \rightarrow \mathrm{GL}(V_{\mathbb{R}}),$$

and a weight space decomposition

$$V_{\mathbb{C}} = \oplus_{p+q=n} V^{p,q}, \text{ where } V^{p,q} = \{v \in V_{\mathbb{C}} : \tilde{\varphi}(z)v = z^p \bar{z}^q v, \}.$$

Definition 3.4. [6] The *Mumford-Tate group* M_{φ} associated to a Hodge structure (V, φ) of weight n is the $\mathbb{Q}$ algebraic closure of

$$\varphi(S^1) \subset \mathrm{SL}(V).$$

For a Hodge structure of even weight $2p$, Hodge classes are defined by $V \cap V_{\mathbb{C}}^{p,p}$.

Theorem 4. [6] *For a Hodge structure (V, φ) of weight n , the Mumford-Tate group M_{φ} is the subgroup of $\mathrm{GL}(V, \mathbb{Q})$, consisting of elements acting as the identity on all the Hodge classes in the tensor products $\oplus V^{\otimes k} \otimes \check{V}^l$ for any pair of integers (k, l) where $n(l - k)$ is even.*

Definition 3.5. [13] An automorphism g of $H^*(X, \mathbb{Q})$ is called a *monodromy operator* if there exists a family $\mathcal{M} \rightarrow B$ (which may depend on g) of irreducible holomorphic symplectic manifolds over quasi-projective scheme B , having X as a fiber over a point $b \in B$, and such that g belongs to the image of $\pi_1(B, b)$ under the monodromy representation. The monodromy group $Mon(X)$ of X is the subgroup, of the automorphism group of the cohomology ring of X , generated by all the monodromy operators.

Denote by $Mon^2(X) \subset GL(H^2(X, \mathbb{Q}))$ the image of $Mon(X)$ in the isometry group of $H^2(X, \mathbb{Z})$. Markman in [13] §4.3 shows that $Mon(X)$ and $Mon^2(X)$ differ only by finite subgroups. Now denote by G_X the identity component of the special orthogonal group associated with the Beauville-Bogomolov form on $H^2(X, \mathbb{Z})$, Proposition 4.1 in [7] shows that the universal cover $\tilde{G}_X \rightarrow G_X$ acts on the cohomology ring of X . Since the cohomology of X is nonzero only in even degrees, this representation passes to G_X .

Let $\mathcal{M}_b$ be a general fiber of the family $\mathcal{M} \rightarrow B$ as in definition 3.5. The work of Deligne in [3] shows that the Mumford-Tate group of the Hodge structure on $H^k(\mathcal{M}_b, \mathbb{Q})$ contains a subgroup of finite index in the monodromy group. This implies that $Mon(X)$ preserves Hodge classes in the tensor products $H^k(\mathcal{M}_b, \mathbb{Q}) \otimes H^l(\mathcal{M}_b, \mathbb{Q})^\vee$ away from a countable union of Zariski closed algebraic subsets, indexed by Hodge classes disappearing under generic deformation.

By the above analysis, the invariants of the G_X representation on $H^*(X, \mathbb{Q})$ are the canonical Hodge classes, i.e. those which remain Hodge under deformations.

Example 3. In table 3, the first row of data shows the number of canonical Hodge classes for $H^{2k}(X, \mathbb{Q})$. For X Kähler deformation equivalent to $S^{[5]}$, there are 2 canonical Hodge classes in $H^{2,2}(X)$, 1 in $H^{3,3}(X)$, 4 in $H^{4,4}(X)$, and 2 in $H^{5,5}(X)$.

3.6. Decomposition of the Cohomology Representation. Denote by G_S the identity component of the special orthogonal group associated with the intersection form on $H^2(S, \mathbb{Z})$.

We can decompose $H^*(X, \mathbb{Q})$ into irreducible representations for the action of G_X . The decomposition

$$H^2(S^{[n]}, \mathbb{Z}) = H^2(S, \mathbb{Z}) \oplus_{\perp} \mathbb{Z}\delta$$

induces an inclusion $G_S \subset G_X$.

[7] provides an explicit method for writing the decomposition:

- (1) Use the isomorphism $H^2(X, \mathbb{Z}) \cong H^2(S, \mathbb{Z}) \oplus_{\perp} \mathbb{Z}\delta$ and compatible maximal torus of G_S and G_X to fix the embedding $G_S \subset G_X$.
- (2) Decompose $H^*(X, \mathbb{Q})$ into the highest-weight representation of G_S by Theorem 2. The highest weight irreducible representation $V_S(\lambda)$ will lie in the summand of the restriction of an irreducible G_X representation $V_X(\lambda)$.
- (3) Repeat step 1 and 2 on $H^*(X, \mathbb{Q})/V_X(\lambda)$.

Now, let $H^{r,n} := H^r(X, \mathbb{Q})$ where X is Kähler deformation equivalent to $S^{[n]}$, and V_λ be the irreducible G_X representation of the highest weight λ ; we get the following computational results

λ	$\dim V_\lambda$	$H^{4,5}$	$H^{6,5}$	$H^{8,5}$	$H^{10,5}$	$H^{6,6}$	$H^{8,6}$	$H^{10,6}$	$H^{12,6}$	...
$(0, 0, 0, \dots)$	1	2	1	4	2	2	5	4	7	...
$(1, 0, 0, \dots)$	23	1	3	3	5	3	4	7	7	...
$(2, 0, 0, \dots)$	275	1	1	3	2	1	4	4	7	...
$(1, 1, 0, \dots)$	253		1	1	2	1	1	3	2	...
$(3, 0, 0, \dots)$	2277		1	1	2	1	1	3	3	...
$(2, 1, 0, \dots)$	4025			1	1		1	2	2	...
$(1, 1, 1, \dots)$	1771								1	...
$(4, 0, 0, \dots)$	14674			1			1	1	2	...
$(3, 1, 0, \dots)$	256795				1			1	1	...
$(2, 2, 0, \dots)$	2193763								1	...
$(5, 0, 0, \dots)$	7804350225				1			1		...
$(4, 1, 0, \dots)$	...							...	...	...
$(6, 0, 0, \dots)$	...								...	...

Table 1: G_X Representations

Here "..." denotes truncated data, and each integer denotes the number of times of V_λ appears in $H^r(X, \mathbb{Q})$. In particular, the first row in the table indicates the number of copies of trivial G_X representation in each $H^r(X, \mathbb{Q})$, and it corresponds to the number of canonical Hodge classes.

$H^2(S, \mathbb{Q})$ corresponds to the standard G_S representation $V_S(1)$. Let $\mathbb{H} := \sum_n H^*(S^{[n]}, \mathbb{Q})[2n]$, which is a bigraded algebra generated by $H^*(S, \mathbb{Q})$ by Theorem 2, and we can decompose $\mathbb{H}_n$ into G_S representations. Let $\mathbb{V}_{n,d} := H^{2n+d}(X, \mathbb{Q})$, the weight k subspace of the G_S representation restricts to the G_X representation on $\mathbb{V}_{k+2n,n}$. By using the same convention as Example 1, we know that

$$\text{Char}_{G_S}(H^k(S^{[n]}, \mathbb{R})) \in \mathbb{Z}[D_1, \dots, D_9]$$

Note that since spin representations don't appear in our representation, we may ignore D^+ and D^- . By the imbedding $G_S \subset G_X$, we obtain

$$(3.1) \quad \text{Char}_{G_S}(H^k(S^{[n]}, \mathbb{R})) = \text{Char}_{G_X}(\mathbb{V}_{k,n})$$

By 3.1, we will not distinguish notations between Char_{G_S} and Char_{G_X} , and will denote both by Char in the following discussion. Let

$$p(z)_n := \sum_{k=-n}^n \text{Char}(\mathbb{V}_{2k+2n,n}) \cdot z^{2k} \in \mathbb{Z}[D_1, \dots, D_9][z, \frac{1}{z}]$$

be the graded character of $\mathbb{H}_n$. Now we take the sum

$$(3.2) \quad p(z, t) := \sum_{n=0}^{\infty} p(z)_n t^n$$

following the grading of each $\mathbb{H}_n$.

Example 4. Consider X as deformation of $S^{[3]}$, and $\mathbb{H}_3 = H^*(X, \mathbb{Q})[6]$. By the analysis in section 3.3 we know that

$$\begin{aligned} H^0(S^{[3]}, \mathbb{Q})[6] &= 1_S \\ H^2(S^{[3]}, \mathbb{Q})[6] &= 1_S \oplus V_S(1) \\ H^4(S^{[3]}, \mathbb{Q})[6] &= 1_S^3 \oplus V_S(1)^2 \oplus V_S(2) \\ H^6(S^{[3]}, \mathbb{Q})[6] &= 1_S^3 \oplus V_S(1)^3 \oplus V_S(2) \oplus V_S(1, 1) \oplus V_S(3) \end{aligned}$$

Thus we have

$$\begin{aligned} p(z)_3 &= \sum_{k=-3}^3 \text{Char}_{G_S}(H^{6+2k}(S^{[3]})) \cdot z^{2k} \\ &= z^{-6} + (1 + D_1)z^{-4} + (3 + 2D_1 + D_1^2 - D_2)z^{-2} \\ &\quad + (3 + 3D_1 + D_1^2 + D_1^3 - 2D_1D_2 + D_3) \\ &\quad + z^6 + (1 + D_1)z^4 + (3 + 2D_1 + D_1^2 - D_2)z^2 \end{aligned}$$

Restricting the above to G_X representation, we obtain

$$\begin{aligned} \mathbb{V}_{3,0} &= H^0(X, \mathbb{Q})[6] = 1_X \\ \mathbb{V}_{3,2} &= H^2(X, \mathbb{Q})[6] = V_X(1) \\ \mathbb{V}_{3,4} &= H^4(X, \mathbb{Q})[6] = 1_X \oplus V_X(1) \oplus V_X(2) \\ \mathbb{V}_{3,6} &= H^6(X, \mathbb{Q})[6] = 1_X \oplus V_X(1) \oplus V_X(1, 1) \oplus V_X(3) \end{aligned}$$

The character formula is

$$\begin{aligned} p(z)_3 &= \sum_{k=-3}^3 \text{Char}_{G_X}(\mathbb{V}_{2k+6,3}) \cdot z^{2k} \\ &= z^{-6} + B_1 z^{-4} + (1 + B_1 + B_1^2 - B_2)z^{-2} + (1 + B_1 + B_2 + B_1^3 - 2B_1B_2 + B_3) \\ &\quad + z^6 + B_1 z^4 + (1 + B_1 + B_1^2 - B_2)z^2 \end{aligned}$$

and the number of canonical Hodge classes of $H^k(X, \mathbb{Q})$ corresponds to the constant term in the coefficient of z^k .

Proposition 5. *The character of the G_X representation is thus*

$$(3.3) \quad p_g(z, t) := \sum_{n=0}^{\infty} p(z)_n t^n = \prod_{m=1}^{\infty} \frac{1}{\det(I_{24} - g t^m)}$$

where $g \in G_X$ is diagonalizable, I_N is a $N \times N$ identity matrix, and $\det(I_{24} - g t^m) = (1 - z^{-2} t^m) \cdot (1 - z^2 t^m) \det(I_{22} - g|_{H^2(S, \mathbb{Q})} t^m)$,

Proof. Recall that $\mathbb{V} = H^*(S, \mathbb{Q})[2] = H^0(S, \mathbb{Q}) + H^2(S, \mathbb{Q}) + H^4(S, \mathbb{Q})$ is bigraded, where $H^0(S, \mathbb{Q})$ and $H^4(S, \mathbb{Q})$ are of weight -2 and $+2$ respectively, and $H^2(S, \mathbb{Q})$ is of weight 0 . For every symmetric power $\text{Sym}^k(\mathbb{V})$, it is sufficient to show the formula holds when g is diagonal. Since G_S acts trivially on H^0 and H^4 , let $u_{-2} \in H^0(S, \mathbb{Q})$ be the eigenvector with eigenvalue 1 and weight -2 , and $u_2 \in H^4(S, \mathbb{Q})$ be eigenvector with eigenvalue 1 and weight 2 . $H^2(S, \mathbb{Q})$ corresponds to the standard G_S representation $V_S(1, 0, \dots, 0)$, and let v_i ($i = 1, \dots, 22$) be its eigenvectors; when i is even, v_i has eigenvalue $t_{\frac{i}{2}}$, and when i is odd, v_i has eigenvalue $t_{\lfloor \frac{i}{2} \rfloor}^{-1}$.

Molien's Formula in [11] indicates that for a representation W of a group G , and given a linear operator $g \in G$, its action on the symmetric algebra $\text{Sym}^\cdot(W)$ has the graded character

$$\sum_{i=0}^{\infty} \text{Char}(\text{Sym}^i(g)) t^i = \frac{1}{\det(I - t g)} .$$

The symmetric algebra on $\mathbb{V}$, denoted by $\text{Sym}^\cdot(\mathbb{V})$, has the form

$$\text{Sym}^\cdot(u_2) \otimes \text{Sym}^\cdot(v_1) \otimes \text{Sym}^\cdot(v_{22}) \otimes \text{Sym}^\cdot(u_{-2}) .$$

Since $v_i \in \mathbb{V}$ is of weight 0 , the bigraded character of $\text{Sym}^\cdot(v_i)$ is

$$\sum_{k=0}^{\infty} (\mu_i z^0)^k \cdot t^k = \frac{1}{1 - \mu_i t} ,$$

where μ_i is the eigenvalue of v_i . Since u_{-2} (resp. u_2) has weight -2 (resp. weight $+2$) in $\mathbb{V}$, we have

$$\begin{aligned} \text{Char}(\text{Sym}^\cdot(u_{-2}))(t) &= \sum_{k=0}^{\infty} (1 \cdot z^{-2})^k t^k = \frac{1}{1 - z^{-2} t} , \text{ and} \\ \text{Char}(\text{Sym}^\cdot(u_2))(t) &= \sum_{k=0}^{\infty} (1 \cdot z^2)^k t^k = \frac{1}{1 - z^2 t} . \end{aligned}$$

Thus, we obtain

$$(3.4) \quad \text{Char}(\text{Sym}^\bullet(\mathbb{V}))(t) = \frac{1}{(1 - z^2 t)(1 - z^{-2} t) \det(I_{22} - g|_{H^2(S, \mathbb{Q})} t)}.$$

By Theorem 2, the graded character $p(z)_n$ of $H^*(X, \mathbb{Q})$ is given by

$$\begin{aligned} p(z)_n t^n &= \text{Char} \left(\sum_{\|\alpha\|=n} \bigotimes_i \text{Sym}^{\alpha_i} \mathbb{V} \right) t^n \\ &= \sum_{\|\alpha\|=n} \prod_i (\text{Char}(\text{Sym}^{\alpha_i} \mathbb{V}) t^{i\alpha_i}). \end{aligned}$$

Note $\text{Char}(\text{Sym}^{\alpha_i} \mathbb{V}) t^{i\alpha_i}$ is the α_i th term in $\text{Char}(\text{Sym}^\bullet(\mathbb{V}))(t^i)$. Then for each $\alpha = (1^{\alpha_1}, 2^{\alpha_2}, \dots, n^{\alpha_n})$ where $\|\alpha\| = n$,

$$\prod_i (\text{Char}(\text{Sym}^{\alpha_i} \mathbb{V}) t^{i\alpha_i})$$

corresponds to the t^n th term in

$$\prod_j (\text{Char}(\text{Sym}^\bullet(\mathbb{V}))(t^j)) ,$$

together with equation 3.4, one can obtain

$$\sum_n p(z)_n t^n = \prod_m \frac{1}{(1 - z^2 t^m)(1 - z^{-2} t^m) \det(I_{22} - g|_{H^2(S, \mathbb{Q})} t^m)}$$

♣

Remark 1. Given a smooth projective complex surface S' , and let $p(S'^{[n]}, z)$ be the Poincaré polynomial $\sum_{i=0}^{4n} \beta_i(S'^{[n]}) z^i$ of $S'^{[n]}$. In [5], Göttsche shows that $\sum_{n=0}^\infty p(S'^{[n]}, z) t^n$ has the expression

$$(3.5) \quad \prod_{m=1}^{\infty} \frac{(1 + z^{2m-1} t^m)^{b_1(S')}(1 + z^{2m+1} t^m)^{b_1(S')}}{(1 - z^{2m-2} t^m)^{b_0(S')}(1 - z^{2m} t^m)^{b_2(S')}(1 - z^{2m+2} t^m)^{b_0(S')}}.$$

If we take g as identity in the equation 3.3, we can obtain the generating series of the Betti numbers of $S^{[n]}$.

Remark 2. We can also apply Weyl measure to obtain the graded generating series $\mathbb{H}^{GS}$ by the following integral

$$\frac{1}{\pi^{11}} \int_0^\pi \cdots \int_0^\pi \left(\left(\prod_{l=1}^{11} (1 - \cos 2(\theta_l)) \right) p(z) \right) d\theta_1 \cdots d\theta_{11}$$

where $e^{i\theta_l} = t_{2l-1}$.

Remark 3. The computation in example 4 is reversible. Explicitly, the number of copies of the highest weight representation $V_X(\lambda)$ appearing in $\mathbb{V}_{n,k}$ corresponds to the coefficient of $\text{Char}(V_X(\lambda))$

4. GENERATING SERIES OF THE CHARACTER OF THE MIDDLE COHOMOLOGY $H^{2n}(X, \mathbb{Q})$

Let $M(q) := \sum_{n=0}^{\infty} \text{Char}(H^{2n}(X, \mathbb{Q})) \cdot q^n$ be the generating series of the character of the representation on the middle cohomology of X . We have

Theorem 6.

$$M(q) = \frac{1 + \sum_{m=1}^{\infty} 2(-1)^m q^{\frac{m(m+1)}{2}}}{\prod_{k=1}^{\infty} \det(I - gq^k)},$$

where $\det(I - gq^k) = (1 - q^k)^2 \cdot \det(I - g|_{H^2(S, \mathbb{Q})})$

Proof. The character of $H^{2n}(X, \mathbb{Q})$ is of weight 0 in equation 3.3. The weight is determined by the representation on $H^0(X, \mathbb{Q})$ and $H^4(X, \mathbb{Q})$. Extract out the weight-controlling part in proposition 5, we obtain

$$\left(\prod_{m=1}^{\infty} \frac{(1 - q^m)^2}{(1 - z^{-2}q^m)(1 - z^2q^m)} \right) \left(\prod_{k=1}^{\infty} \det(I - gq^k) \right)^{-1}$$

Let $a_k(q) := \sum_{l=1}^{\infty} (-1)^{l+1} q^{\frac{l(l+2k-1)}{2}}$, the z^{2k} th coefficient is $a_k(q) - a_{k+1}(q)$. In particular, z^0 th term is

$$a_0(q) - a_1(q) = 1 + \sum_{l=1}^{\infty} 2(-1)^l q^{\frac{l(l+1)}{2}}$$

which gives the formula. ♣

Corollary 7.

$$\sum_{n=0}^{\infty} \dim(H^{2n+2k}(X, \mathbb{Q})) q^n = \frac{q}{\Delta(q)} (a_k(q) - a_{k+1}(q))$$

where $\Delta(q) = q \prod_k (1 - q^k)^{24}$ is a cusp form of weight 12 for $SL_2(\mathbb{Z})$, and $a_k(q) := \sum_{l=1}^{\infty} (-1)^{l+1} q^{\frac{l(l+2k-1)}{2}}$.

Remark 4. [5] shows that the generating series for the Euler numbers of $S^{[n]}$ is

$$\sum_{n=0}^{\infty} e(S^{[n]}) q^n = \frac{q}{\Delta(q)}.$$

According to [16] and remark 3.7 of Appendix in [10] we have

$$\Delta(\tau) = 4096\epsilon(\delta^2 - \epsilon)^2$$

where ϵ and δ are modular forms for $\Gamma_0(2)$ of weights 4 and 2 with the following forms:

$$\begin{aligned}\epsilon &= \sum_{n=1}^{\infty} \left(\sum_{d|n, \frac{n}{d} \text{ odd}} d^3 \right) q^n \\ \delta &= -\frac{1}{8} - 3 \sum_{n=1}^{\infty} \left(\sum_{d|n, d \text{ odd}} d \right) q^n\end{aligned}$$

It is interesting that the modular form Δ appears many times in computations related to the cohomology rings of $S^{[n]}$.

5. APPENDIX: TABLE OF NONSTABLE G_X REPRESENTATIONS

λ	$H^{8,7}$	$H^{10,7}$	$H^{12,7}$	$H^{14,7}$	$H^{8,8}$	$H^{10,8}$	$H^{12,8}$	$H^{14,8}$	$H^{16,8}$	$H^{10,9}$	$H^{12,9}$	$H^{14,9}$	$H^{16,9}$	$H^{18,9}$	...
(0, 0, 0, ..)	5	5	10	7	6	6	13	12	18	6	15	15	25	21	...
(1, 0, 0, ..)	5	9	11	14	5	10	14	21	21	11	16	27	33	39	...
(2, 0, 0, ..)	4	5	10	9	4	6	13	15	21	6	14	19	31	30	...
(1, 1, 0, ..)	1	4	4	7	1	4	5	10	9	4	6	13	15	21	...
(3, 0, 0, ..)	1	4	5	7	1	4	6	11	11	4	7	14	18	24	...
(2, 1, 0, ..)	1	2	4	5	1	2	5	8	10	2	5	10	16	18	...
(1, 1, 1, ..)			1				1	1	2		1	1	3	3	...
(4, 0, 0, ..)	1	1	3	3	1	1	4	5	8	1	4	6	12	11	...
(3, 1, 0, ..)		1	2	3		1	2	5	5	1	2	6	9	13	...
(2, 2, 0, ..)			1				1	1	3		1	1	4	3	...
(2, 1, 1, ..)				1				1	1			1	2	3	...
(5, 0, 0, ..)		1	1	2		1	1	3	3	1	1	4	5	8	...
(4, 1, 0, ..)			1	1			1	2	3		1	2	5	6	...
(3, 2, 0, ..)				1				1	1			1	2	3	...
(3, 1, 1, ..)									1				1	1	...
(2, 2, 1, ..)														1	...
(6, 0, 0, ..)			1				1	1	2		1	1	3	3	...
(5, 1, 0, ..)				1				1	1			1	2	3	...
(4, 2, 0, ..)									1				1	1	...
(4, 1, 1, ..)														1	...
(3, 3, 0, ..)														1	...
(7, 0, 0, ..)				1				1				1	1	2	...
(6, 1, 0, ..)									1				1	1	...
(5, 2, 0, ..)														1	...
(8, 0, 0, ..)									1				1		...
(7, 1, 0, ..)														1	...
(9, 0, 0, ..)														1	...

REFERENCES

- [1] B. Bakker and A. Jorza. Lagrangian hyperplanes in holomorphic symplectic varieties. *arXiv preprint arXiv:1111.0047*, 2011.
- [2] Arnaud Beauville. Variétés Kähleriennes dont la première classe de Chern est nulle. *J. Differential Geom.*, 18(4):755–782 (1984), 1983.
- [3] Pierre Deligne. La conjecture de Weil pour les surfaces $K3$. *Invent. Math.*, 15:206–226, 1972.
- [4] William Fulton and Joe Harris. *Representation theory, a first course*, volume 129 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991.
- [5] Lothar Göttsche. The Betti numbers of the Hilbert scheme of points on a smooth projective surface. *Math. Ann.*, 286(1-3):193–207, 1990.
- [6] Mark Green, Phillip Griffiths, and Matt Kerr. *Mumford-Tate groups and domains*, volume 183 of *Annals of Mathematics Studies*. Princeton University Press, Princeton, NJ, 2012.
- [7] David Harvey, Brendan Hassett, and Yuri Tschinkel. Characterizing projective spaces on deformations of Hilbert schemes of $K3$ surfaces. *Comm. Pure Appl. Math.*, 65(2):264–286, 2012.
- [8] Brendan Hassett and Yuri Tschinkel. Moving and ample cones of holomorphic symplectic fourfolds. *Geom. Funct. Anal.*, 19(4):1065–1080, 2009.
- [9] Brendan Hassett and Yuri Tschinkel. Intersection numbers of extremal rays on holomorphic symplectic varieties. *Asian J. Math.*, 14(3):303–322, 2010.
- [10] Friedrich Hirzebruch, Thomas Berger, and Rainer Jung. *Manifolds and modular forms*. Aspects of Mathematics, E20. Friedr. Vieweg & Sohn, Braunschweig, 1992. With appendices by Nils-Peter Skoruppa and by Paul Baum.
- [11] Stavros Kousidis. A closed character formula for symmetric powers of irreducible representations. In *22nd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2010)*, Discrete Math. Theor. Comput. Sci. Proc., AN, pages 833–844. Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2010.
- [12] Manfred Lehn and Christoph Sorger. The cup product of Hilbert schemes for $K3$ surfaces. *Invent. Math.*, 152(2):305–329, 2003.
- [13] Eyal Markman. On the monodromy of moduli spaces of sheaves on $K3$ surfaces. *J. Algebraic Geom.*, 17(1):29–99, 2008.
- [14] Hiraku Nakajima. Heisenberg algebra and Hilbert schemes of points on projective surfaces. *Ann. of Math. (2)*, 145(2):379–388, 1997.
- [15] Mikhail Sergeevic Verbitsky. *Cohomology of compact hyperkaehler manifolds*. ProQuest LLC, Ann Arbor, MI, 1995. Thesis (Ph.D.)—Harvard University.
- [16] Don Zagier. Note on the Landweber-Stong elliptic genus. In *Elliptic curves and modular forms in algebraic topology (Princeton, NJ, 1986)*, volume 1326 of *Lecture Notes in Math.*, pages 216–224. Springer, Berlin, 1988.

DEPARTMENT OF MATHEMATICS, RICE UNIVERSITY, 6100 MAIN STREET,
HOUSTON, TX 77005, U.S.A.

E-mail address: Letao.zhang@rice.edu